


In the Claims:

Claims

1. **(currently amended):** A process for the preparation of secondary nitroxide radicals from their corresponding secondary amines by oxidation with an organic peracid, comprising the steps
a) adding to a reaction vessel a secondary amine, optionally together with an organic solvent, and in one batch a base selected from the group consisting of alkali metal, alkaline earth metal or ammonium bicarbonates and alkaline earth metal or ammonium carbonates or mixtures thereof in the form of a solid together with water or as an aqueous slurry;
b) dosing a peracid under stirring to the reaction mixture in an amount of ~~1.0 to 2.5~~ 1.0 to 2.5 mol per mol of secondary amine; and
c) isolating the organic phase.
2. **(original):** A process according to claim 1 wherein the organic solvent optionally used in step a) is immiscible with water.
3. **(original):** A process according to claim 1 wherein the peracid is peracetic acid.
4. **(original):** A process according to claim 1 wherein the amount of water added in step a) is sufficient to dissolve the organic acid salt formed in the neutralization reaction between the organic acid and the bicarbonate or carbonate.
5. **(original):** A process according to claim 1 wherein the base is sodium or potassium bicarbonate, calcium or magnesium carbonate or dolomite.
6. **(original):** A process according to claim 1 wherein the base is added in an amount of from 0.1 to 1.5 equivalents base per 1 equivalent of all acids present.
7. **(original):** A process according to claim 1 wherein the reaction temperature is between 0° C and 40° C.
8. **(original):** A process according to claim 1 wherein the dosage of the peracid is carried out from 10 minutes to 5 hours.


9. (original): A process according to claim 1 wherein the nitroxide radical is of formula (I)



R_p and R_q are independently tertiary bound C_4 - C_{28} alkyl groups or C_3 - C_{17} secondary bound alkyl groups which are unsubstituted or substituted by one or more electron withdrawing groups or by phenyl; or

R_p and R_q together form a 5, 6 or 7 membered heterocyclic ring which is substituted at least by 4 C_1 - C_4 alkyl groups and which may be interrupted by a further nitrogen or oxygen atom.

10. (original): A process according to claim 1 wherein the nitroxide radical contains a structural



G_1 , G_2 , G_3 , G_4 are independently C_1 - C_6 alkyl or G_1 and G_2 or G_3 and G_4 , or G_1 and G_2 and G_3 and G_4 together form a C_5 - C_{12} cycloalkyl group;

*denotes a valence; and

G_5 , G_6 independently are H or C_1 - C_6 alkyl.